

Evaluation of genetic diversity among Iranian pomegranate (*Punica granatum* L.) cultivars, using ISSR and RAPD markers

Majid Talebi Bedaf^{1*}, Masoud Bahar¹, Bahram Sharifnabi¹ and Ahad Yamchi²

¹ Department of Agricultural Biotechnology, College of Agriculture, Isfahan University of Technology, Isfahan, Iran, 84156-83111.

² Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran

Abstract

Considering the high level of morphological diversity in Iranian pomegranate cultivars, comparison of genetic variation among 24 pomegranate cultivars was evaluated using random amplified polymorphic DNA (RAPD) and inter simple sequence repeat (ISSR) markers. RAPD primers amplified 131 DNA fragments among which 29 were polymorphic (22.14%) and ISSR markers produced 173 amplification products, out of which 64 were polymorphic (37%). Mean PIC (polymorphic information content) was 0.128 for RAPD and 0.163 for ISSR. The results suggested that the ISSR markers produced much better reproducible bands and were more efficient in grouping cultivars. Pairwise similarity index values ranged from 0.353 to 1.0 (RAPD), 0.291 to 0.930 (ISSR) and mean similarity index values of 0.604 and 0.674 for RAPD and ISSR, respectively. The analysis of molecular variance (AMOVA) for RAPD and ISSR data showed no significant differences among the geographical regions and juice acidity of the used cultivars ($P>0.05$) indicated that genetic and geographic distances were not correlated.

Key words: *Punica granatum*, genetic diversity, pomegranate, RAPD, ISSR

Introduction

Pomegranate (*Punica granatum* L.) belongs to Punicaceae family and is an important fruit tree of tropical and subtropical regions of the world which is valued highly for its delicious edible fruits. In addition, the tree is also cultivated for its pharmaceutical and ornamental usages (Levin, 1994). The pomegranate tree has a wide geographical distribution that spreads from Iran to the Himalayas in northern India, and has been cultivated since ancient times throughout the Mediterranean regions of Asia, Africa and Europe (Levin, 1994). Pomegranate may be classified according to the acidity of its fruit into sour, sour-sweet or sweet.

Development of highly reliable and discriminatory methods have become increasingly important to plant breeders for identifying cultivars and to those in the nursery industry who need sensitive tools to differentiate and identify cultivars for plant patent protection

* Corresponding Author: mtalebi@cc.iut.ac.ir

(Wunsch and Hormaza, 2002). In the past, cultivars were identified primarily based on horticultural, morphological and physiological descriptions. In most cases, the descriptions and measurements varied considerably due to environmental fluctuation and differences human judgment. Differences in DNA sequence among individuals could be detected by different methods. Almost all kind of DNA markers can be used for fingerprinting fruit tree species (Wunsch and Hormaza, 2002). The Random Amplified Polymorphic DNA (RAPD) technique (Williams *et al.*, 1990) based on the Polymerase Chain Reaction (PCR) has been used to detect polymorphism in some species (Williams *et al.*, 1990). Thus, the RAPD technique can generate polymorphisms between very closely related genotypes. Since 1990, RAPD markers have been successfully used to identify cultivars and/or clones of various plant species (Belaj *et al.*, 2001; Besnard *et al.*, 2001; Claros *et al.*, 2001; Ozden-Tokatli *et al.*, 2010; Takeda *et al.*, 1998).

Also, Inter-simple sequence repeat (ISSR) amplification is a technique which can rapidly differentiate closely related individuals (Zietkiewicz *et al.*, 1994). ISSR markers involve PCR amplification of DNA using a single primer composed of a microsatellite sequence such as (CA)₈ anchored at the 3' or 5' end by 2-4 arbitrary, often degenerate nucleotides. The sequences of repeats and anchored nucleotides are randomly selected. Coupled with the separation of amplifications products on a polyacrylamide gel, ISSR amplification can reveal a higher number of fragments per primer than RAPD. ISSR markers have been used for cultivar identification and for genetic relationship studies in various plant species (Awasthi *et al.*, 2004; Martin and Sanchez-Yelamo, 2000; Weiguo *et al.*, 2007).

Although a wide range of morphological and physiological characters show variabilities in the pomegranate, molecular studies of the pomegranate have been restricted to examinations of RAPD (Dorgac *et al.*, 2008; Zamani *et al.*, 2007; Sarkhosh *et al.*, 2009), ISSR (Talebi Bedaf *et al.*, 2005), AFLP (Jbir *et al.*, 2008; Rahimi *et al.*, 2006) and SSR (Koochi-Dehkordi *et al.*, 2007; Ebrahimi *et al.*, 2010; Pirseyedi *et al.*, 2010) to investigate the population dynamics of economically important cultivars. To meet various breeding programs and to conserve the existing genetic resources of pomegranate, the objectives of this study were to assess the levels of polymorphisms detected by RAPD and ISSR markers, comparison of information content the marker systems and using them to identify 24 Iranian pomegranate cultivars.

Materials and Methods

Plant materials and DNA extraction

Twenty four cultivars of *P. granatum* were collected from Agricultural Research Center of Yazd province, Yazd, Iran. The selection of cultivars was based on the acidity of fruits and the morphological characteristics, such as color and shape of fruit (Table 1).

Total DNA was extracted from young leaves following the CTAB (Hexadecyltrimethylammonium bromide) method described by Murray and Thompson (1980) with modifications. The purified total DNA was quantified by agarose gel electrophoresis and its quality was verified by spectrophotometry. DNA samples were diluted to 25ng/μl and stored at -20 °C.

Table 1: Iranian pomegranate genotypes included in the study

No.	Cultivar codes	Cultivar names	Acidity	Origin
1	PGpsy	Poust syah yazdi	Sweet	Yazd
2	PGgsh	Goroch shahvar	Sweet	Yazd
3	PGtl	Tab va larz	Sweet	Yazd
4	PGbl	Bihasteh ladiz	Sweet	Systan va Balouchestan
5	PGas	Asali sarvestan	Sweet	Fars
6	PGghr	Golabi hasteh riz	Sour-sweet	Systan va Balouchestan
7	PGna	Nabati ardakan	Sour-sweet	Yazd
8	PGdhv	Dokhtar hamoumi varamin	Sour-sweet	Tehran
9	PGggn	Galu gandeh neiriz	Sour-sweet	Fars
10	PGakh	Amaneh khatouni	Sour-sweet	Yazd
11	PGtg	Togh gardan	Sour-sweet	Yazd
12	PGda	Dom anbarouti	Sour	Khorasan
13	PGpa	Panjeh arous khafr	Sour	Fars
14	PGapgh	Ardestani poust ghermez	Sour	Esfahan
15	PGvns	Vahshi narak sarvestan	Sour	Fars
16	PGds	Dabbei sarjangal	Sour	Kerman
17	PGgf	Golnar fars	Ornamental	Fars
18	PGkgs	Kaleh gavi sangan	Sour-sweet	Systan va Balouchestan
19	PGsh	Shahvar shirin	Sweet	Yazd
20	PGhm	Hasibi mehriz	Sour-sweet	Yazd
21	PGbs	Bihasteh sangan	Sweet	Systan va Balouchestan
22	PGaps	Ardestani poust sefid	Sweet	Esfahan
23	PGtmz	Torsh mamuli zabol	Sour	Systan va Balouchestan
24	PGsb	Sabi bam	Sour	Kerman

RAPD assay

One hundred-four 10mer oligonucleotide primers among sets A, B, C, AC, AD, AE, P and S (Operon technologies, Inc, USA), AJ, MG and UBC (Roche Molecular Biochemicals, Germany) were used as single primers for DNA amplification and 13 of them were selected based on clear and reproducible banding patterns. The PCR was performed in a Genius (FGENO5TD) Thermal Cycler, in a 25µl volume containing of 2.5µl of 1X reaction buffer [100mM Tris-HCl, 15mM MgCl₂, 500mM KCl, pH 8.3 (20°C)], 0.5mM MgCl₂, 200µM each of dNTPs (Roche, Germany), 0.4µM of 10mer primer, 0.75 units of *Taq* DNA polymerase (Roche, Germany) and 100ng of template DNA, overlaid with 25µl of sterile mineral oil. The amplification condition was: initial step of denaturation at 94°C for 2 min followed by 40 cycles of denaturation at 92°C for 1 min, primer annealing at 35°C for 1 min and extension at 72°C for 2 min, followed by an extended elongation step at 72°C for 5 min. The amplification products were analysed on 1.2% MP agarose gel (Roche, Germany) in 1X TBE buffer running at 60 volts for three hours and stained in ethidium bromide (0.5mg/ml) and visualized under UV light and photographed. The DNA size marker used was 1Kb ladder (Life technologies).

ISSR assay

A total of 15 primers were tested to amplify DNA from which six primers with considerable polymorphism and reproducibility were selected for further analysis (Table 2). PCR were performed in 15 µl volume consisted of 1X PCR buffer, 2mM MgCl₂, 200 µM each of dNTPs, 1 µM primer, 1 U of *Taq* DNA polymerase (Roche, Germany), 2% formamide and 25 ng of template DNA. Each reaction mixture was overlaid with 25 µl of sterile mineral oil. Amplification was performed in a Genius (FGENO5TD) Thermal Cycler under the following conditions: 4 min at 94°C for 1 cycle, followed by 30 s at 94°C, 45 s at

52°C, and 2 min at 72°C for 30 cycle, and 5 min at 72°C for a final extension. Amplification products were separated on 6% denaturing polyacrylamide gels containing 7 M Urea and 1X TBE buffer in Biometra sequencing gel (S2 model). Amplified DNA segments were detected using silver staining (Bassam *et al.*, 1991).

Data analysis

A marker index was calculated for the RAPD and ISSR markers to characterize the capacity of each primer to detect polymorphic loci among the cultivars. As such, the marker index was the sum of the polymorphism information content (PIC) values for all the selected markers produced by a particular primer. The PIC value was calculated using the formula $PIC=1-\sum p_i^2$, where P_i is the frequency of the i allele (Smith *et al.*, 1997).

Only reproducible and clear bands in the replications were considered as potential polymorphic markers. The data obtained by scoring the presence (1) or absence (0) of amplified fragments from the RAPD, ISSR and also collective of marker profiles were subjected to the construction of a similarity matrix using Jaccard's (Jaccard, 1908) coefficients of similarity. The matrices were then used for a cluster analysis. Sequential agglomerative hierarchical nonoverlapping (SAHN) clustering was performed using the unweighted pair group method with arithmetic averages (UPGMA), and then, the results were summarized as dendrograms using NTSYSpc software 2.02 (Raholf, 1998).

The pertinency of the dendrograms to the original similarity matrix was calculated by computing the cophenetic values (rcoph) using the cophenetic (COPH) and matrix comparison (MXCOMP) modules of NTSYSpc.

Finally, the frequency of occurrence of each marker in each cultivar was computed, to render a matrix of 24 cultivars by RAPD and ISSR markers. These matrices were afterward subjected to principal component analysis (PCA).

Analysis of molecular variance (AMOVA) was performed to estimate variance components for RAPD and ISSR data and partitioning the variation into within and among local regions and acidity of cultivars, using Arlequin 3.1 software (Excoffier *et al.*, 2005).

Results

One hundred-four 10 mer RAPD primers were screened and among which 13 were chosen for their clear and reproducible band patterns (Table 2). The thirteen selected primers generated 131 RAPD fragments, an average of 10.08 bands per primer. The size of the amplified products ranged from 400 to 3,000 bp. The total number of polymorphic markers and percentage of polymorphism were 29 and 22.14%, respectively (Table 2). Primers OPAD02 and MG16 amplified maximum number of polymorphic bands. Primers OPAD02 and OPAE10 put out the highest level of distinguishable polymorphism. The PIC values, a reflection of the allele diversity and frequency among the cultivars, were not uniform with respect to for all the RAPD loci tested. The PIC values ranged from 0.012 (MG01) to 0.373 (OPAD02) with a mean of 0.128. The result showed that the minimum similarity (0.353) existed in the two local cultivars "Dom anbarouti" and "Poust syah yazdi" and the maximum similarity (1.00) occurred in the cultivars "Tab va larz" and "Bihasteh ladiz". The mean similarity index was 0.604. The data obtained from RAPD analysis of 24 pomegranate cultivars was subjected to UPGMA analysis. The cophenetic correlation coefficient (0.91) indicated little distortion between the original similarity values from the similarity matrix and the values used to construct the dendrogram. A cluster analysis was performed based on Jaccard's similarity coefficient matrices, calculated from the RAPD

markers. At the similarity of 59%, twenty cultivars were categorized in one group and the four remaining cultivars were placed in the separate groups (Figure 1a).




Figure 1: UPGMA dendrogram showing relations among Iranian pomegranate cultivars using RAPD (a) and ISSR (b) data and Jaccard's similarity coefficient.

Eighteen ISSR primers were initially tested using pomegranate DNA as single or combined. Three primers as single and three as combined were based on $(AG)_n$, $(GT)_n$, $(GA)_n$, $(AC)_n$, $(CT)_n$ or $(CA)_n$ repeats, each anchored by various nucleotides used in this study as polymorphic primers (Table 2). The six selected primers generated 173 fragments, an average of 28.83 bands per primer. The size of the amplified products ranged from 80 to 3,000 bp with the scoreable region being from 100 to 2,000 bp accordingly, and the total number of polymorphic markers and percentage of polymorphism were 64 and 36.99%, respectively (Table 2). In the case of the ISSR analysis, the mean PIC value was 0.163, and the lowest and highest PIC values were 0.099 (ISSR5 and ISSR6) and 0.257 (ISSR11), respectively. The similarity coefficients for 24 pomegranate cultivars based on the ISSR fragments ranged from 0.291 ("Dabbei sarjangal" and "Poust syah yazdi") to 0.930 ("Dom anbarouti" and "Golabi hasteh riz"). The mean similarity index was 0.674. Twenty four cultivars were clustered in five distinct groups at the similarity level of 65%, 19 of which were placed in one group (Figure 1b).

Table 2: RAPD and ISSR primers successfully used in this study and the number of total and polymorphic bands amplified in pomegranate cultivars.

Marker name	Primer name	Sequence (3'-5')	Total band number	Number of polymorphic bands	Polymorphic bands (%)	PIC value	
RAPD	OPAC11	CCTGGGTCAG	12	3	25	0.188	
	OPAD02	CTGAACCGCT	8	5	62.5	0.373	
	OPAD04	GTAGGCCTCA	8	2	25	0.246	
	OPAD13	GGTTCCTCTG	5	2	40	0.172	
	OPAD15	TTTGCCCCGT	6	1	16.67	0.014	
	OPAD16	AACGGGCGTC	10	2	20	0.110	
	OPAE10	CTGAAGCGCA	11	2	18.18	0.081	
	OPB10	CTGCTGGGAC	12	1	8.33	0.013	
	OPP02	TCGGCACGCA	10	2	20	0.135	
	OPP14	CCAGCCGAAC	12	2	16.67	0.096	
	MG01	AGCGCCGACG	14	2	14.29	0.012	
	MG11	AGGAGCTGCC	16	2	12.5	0.024	
	MG16	GAAGAACCGC	7	3	42.86	0.200	
	Total			131	29		
	Mean			10.08	2.23	22.14	0.128
	ISSR	LK7	5'-CCA(CT) ₈ -3'	32	10	31.25	0.256
		ISSR5	5'-CCA(AG) ₈ T-3'	25	8	32	0.099
ISSR6		5'-(GA) ₈ C-3'					
ISSR10		5'-(GT) ₈ A-3'	38	11	28.94	0.106	
ISSR12		5'-(AG) ₈ YT-3'					
ISSR11		5'-(AG) ₈ YT-3'	24	6	25	0.156	
ISSR12		5'-(GA) ₈ YT-3'					
ISSR11		5'-(AG) ₈ YT-3'	26	15	57.69	0.257	
ISSR12		5'-(GA) ₈ YT-3'	28	14	50	0.103	
Total				173	64		
Mean				28.83	10.67	36.99	0.163

The matrices for RAPD and ISSR markers were also compared using Mantel's test (Mantel, 1967) for matrix correspondence. The correlation between the matrices of cophenetic values relating to the dendrograms based on RAPD and ISSR data was very low ($r=0.02$).

The relationships among cultivars were initially defined by the first three principal vectors of the PCA, which together accounted for 65% (RAPD), 73% (ISSR) and 66% (RAPD+ISSR) of the total variation at the molecular level (data not shown).




Figure 2: UPGMA dendrogram of 24 pomegranate cultivars based on RAPD and ISSR pulled data and Jaccard's similarity coefficient.

The similarity coefficients of 24 pomegranate cultivars based on 29 RAPD and 64 ISSR markers ranged from 0.338 ("Dabbei sarjangal" and "Poust syah yazdi") to 0.932 ("Bihasteh ladiz" and "Tab va larz"), and accordingly the mean similarity index value of the combined RAPD and ISSR was 0.655. A cluster analysis performed based on combination of the data for both markers, separated the cultivars into two distinct clusters. The first cluster included only two cultivars, whereas the second cluster was further divided into three subclusters. Among these subclusters, two local cultivars ("Poust syah yazdi" and "Goroch shahvar") were separated from other cultivars (Figure 2).

AMOVA for RAPD and ISSR data indicated that there were no significant differences among the geographical regions and juice acidity of the used cultivars ($P > 0.05$).

Discussion

Among markers, RAPD and ISSR are simple, which provide a quick screen for DNA polymorphism and very small amounts of DNA are required. In addition, information on template DNA sequence is not necessary. However, with respect to RAPD markers problems of reproducibility are reported (Muthusamy *et al.*, 2008). In order to assure reproducibility, optimization of PCR reaction and also its repetition is essential. In this study, each RAPD analysis was repeated in separate experiments at least twice, and only reproducible markers were considered. ISSR primers consist of 17-19 nucleotides and optimization of annealing temperature is important. The choice of annealing temperature for further ISSR analysis is based on the complexity and reproducibility of banding patterns. The primers that were based on (AT)_n or (TA)_n repeats amplified no products at all. Possibly, this indicates that the pomegranate genome lacks, or else has very few of these two microsatellites, although Wang *et al.*, 1994, reported that (AT)_n was the most abundant microsatellite in plant nuclear genomes. Alternatively, lack of amplification products may be due to the self-complementary nature of (AT)_n or (TA)_n primers. In this study, when optimal conditions for PCR had been determined, reproducible patterns were obtained for both RAPD and ISSR assays.

In general, among the set of accessions investigated, the efficiency of a molecular marker technique depends on the amount of polymorphism it can detect. In our study, ISSR fingerprinting was more efficient than the RAPD assay; it detected 37% polymorphic DNA markers among the 24 cultivars analyzed, compared with 22.14% for RAPD fingerprinting. Similar results were obtained for several other plant species (Galvan *et al.*, 2003; Nkongolo *et al.*, 2005; Qian *et al.*, 2001; Raina *et al.*, 2001). However, Fang and Roose (1997) showed that RAPD had a higher level of variation in *Citrus* spp. than ISSR, and Metais *et al.*, (2000) demonstrated that the two techniques produced similar levels of polymorphism in *Phaseolus vulgaris*. The correlation between the matrices of cophenetic values for the dendrograms based on RAPD and ISSR data was also very low ($r=0.02$). It is probably due to the nature of different marker systems. RAPD markers cover the entire genome, revealing length polymorphisms in coding or noncoding and repeated or single-copy sequences (Williams *et al.*, 1990), whereas, the origin of the amplification products in ISSR is known to be from the sequences between the two microsatellite sites (Zietkiewicz *et al.*, 1994).

Three first principle eigen vectors of the PCA, which together accounted for RAPD, ISSR and combined data showed high total variation at the molecular level, indicating the suitability of the RAPD and ISSR approaches for genetic clustering.

Based on the pairwise analysis of the amplification products which were obtained with

the 13 tested RAPD primers, all the tested pomegranate cultivars showed a very high similarity values. Different relationships were observed between various cultivars. "Bihasteh sangan", "Dom anbarouti" and "Golabi hasteh riz" were quite distinct from the rest of cultivars. "Tab va larz" and "Bihasteh ladiz" were clustered together and in the studied cultivars showed the highest average similarity value (similarity coefficient of 1.0) which indicates that although there are some morphological differences in fruit characteristics, these cultivars may probably be mutants of each other (Sarkhosh *et al.*, 2009). All the remaining cultivars showed very limited differences, but sufficient to distinguish the different cultivars.

Among the 24 cultivars analyzed with ISSR marker, four main groups were recognized by UPGMA based on Jaccard's similarity coefficient (Figure 1b). The first group contained "Poust syah yazdi", the second group included "Goroch shahvar" and "Vahshi narak sarvestan", the third group consisted of "Dabbei sarjangal" and "Kaleh gavi sangan", while all remaining cultivars formed the fourth group. "Vahshi narak sarvestan", "Dabbei sarjangal" and "Kaleh gavi sangan" were quite distinct from the rest of cultivars and were readily separated from other cultivars. These cultivars are wild and it seems that they differ from other cultivars, morphologically e.g. fruit size, fruit color, seed color and the taste. "Dom anbarouti" and "Golabi hasteh riz" grouped and showed the highest average similarity value among the studied cultivars. "Poust syah yazdi" cultivars that have a black bark as a distinguishable marker from the other cultivars, was separate in both RAPD and ISSR analysis. In total, ISSR analysis, was more efficient than RAPD analysis.

A close genetic similarity was found in some of the cultivars analyzed as shown by high values of similarity index. Also, the similarities detected with ISSRs are greater than the similarities measured according to RAPD data. Fernandez *et al.*, (2002) and Muthusamy *et al.*, (2008) have studied barley cultivars and rice bean (*Vigna umbellata*) landraces, respectively, and they also found higher similarity index by ISSRs than by RAPDs.

Observation of no significant difference among the geographical regions based on AMOVA for RAPD and ISSR data and the clustering pattern of cultivars revealed that there are no correlation between genetic diversity and geographic distances. In the study of RAPD profiles in Iranian pomegranates by Sarkhosh *et al.*, (2009) and of AFLP profiles in Tunisian pomegranates by Jbir *et al.*, (2008), the authors could not detect any correlation between provenance of the accessions and similarity or otherwise in the PCR profiles. This can be due to the exchange of plant materials across the regions during the history of pomegranate cultivation.

In conclusion, both RAPD and ISSR are methods useful for revealing molecular relationships among pomegranate cultivars and ISSR markers exhibited higher levels of polymorphisms than RAPD. Relationships among these pomegranate cultivars revealed by ISSR markers were not generally in agreement grouping showed by RAPD markers.

Acknowledgments

We are grateful to Seyed Ziaaddin Tabatabaei-Ardakani from Agricultural Research Center of Yazd province for providing pomegranate cultivars and all of the colleagues in biotechnology center of IUT.

References

- Awasthi, A. K., Nagaraja, G. M., Naik, G. V. S., Thangavelu, K. and Nagaraju, J. (2004) Genetic diversity and relationships in mulberry (genus *Morus*) as revealed by RAPD and ISSR marker assays. BMC Genetics 5: 1.

- Bassam, B. J., Caetano-Anolles, G., Gresshoff, P. M. (1991) Fast and sensitive silver staining of DNA in polyacrylamide gels. *Analytical Biochemistry* 196: 80-83.
- Belaj, A., Trujillo, I., Rosa, R. D. L. and Rallo, L. (2001) Polymorphism and discrimination capacity of randomly amplified polymorphic markers in an olive germplasm bank. *Journal of American Society of Horticultural Science* 126: 64-71.
- Besnard, G., Baradat, P. and Berville, A. (2001) Genetic relationships in the olive (*Olea europaea* L.) reflect multilocal selection of cultivars. *Theoretical and Applied Genetics* 102: 251-258.
- Claros, M. G., Crespillo, R., Aguilar, M. L. and Canovas, F. M. (2001) DNA fingerprinting and classification of geographical related genotypes of olive tree (*Olea europaea* L.). *Euphytica* 116: 131-142.
- Durgac, C., Ozgen, M., Simsek, O., Kacar, Y. A., Kiyga, Y., Celebi, S., Gunduz, K. and Serce S (2008) Molecular and pomological diversity among pomegranate (*Punica granatum* L.) cultivars in Eastern Mediterranean region of Turkey. *African Journal of Biotechnology* 7: 1294-1301.
- Ebrahimi, S., Sayed-Tabatabaei, B. E. and Sharifnabi, B. (2010) Microsatellite isolation and characterization in pomegranate (*Punica granatum* L.). *Iranian Journal of Biotechnology* 8: 156-163
- Excoffier, L., Laval, G. and Schneider, S. (2005) Arlequin ver. 3.0: An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1: 47-50.
- Fang, D. Q. and Roose, M. L. (1997) Identification of closely related citrus cultivars with inter-simple sequence repeat markers. *Theoretical and Applied Genetics* 95: 408-417.
- Fernandez, M.E., Figueiras, A.M. and Benito, C. (2002) The use of ISSR and RAPD markers for detecting DNA polymorphism, genotype identification and genetic diversity among barley cultivars with known origin. *Theoretical and Applied Genetics* 104: 845-851.
- Galvan, M. Z., Bornet, B., Balatti, P. A. and Branchard, M. (2003) Inter simple sequence repeat (ISSR) marker as a tool for the assessment of both genetic diversity and gene pool origin in common bean (*Phaseolus vulgaris* L.). *Euphytica* 132: 297-301.
- Jaccard, P. (1908) Nouvelles recherches sur la distribution florale. *Bulletin de la Societe. Vaudoise des Sciences Naturelles* 44: 223-270.
- Jbir, R., Hasnaoui, N., Mars, M., Marrakchi, M. and Trifi, M. (2008) Characterization of Tunisian pomegranate (*Punica granatum*) cultivars using amplified fragment length polymorphism analysis. *Scientia Horticulturae* 115: 231-237.
- Koohi-Dehkordi, M., Sayed-Tabatabaei, B. E., Yamchi, A. and Danesh-shahraki, A. (2007) Microsatellites markers in pomegranate. *Acta Horticulture* 760: 179-183.
- Levin, G.M. (1994) Pomegranate (*Punica granatum*) plant genetic resources in Turkmenistan. *Plant Genetic Resource Newsletters* 97: 31-36.
- Mantel, N. (1967) The detection of disease clustering and a generalized regression approach. *Cancer Research* 27: 209-220.
- Martin, J. P. and Sanchez-Yelamo, M. D. (2000) Genetic relationship among species of the genus *Diplotaxis* (Brassicaceae) using intersimple sequence repeat markers. *Theoretical and Applied Genetics* 101: 1234-1241.
- Metais, I., Aubry, C., Hamon, B., Jalouzot, R. and Peltier, D. (2000) Description and analysis of genetic diversity between commercial bean lines (*Phaseolus vulgaris*, L.). *Theoretical and Applied Genetics* 101: 1207-1214.
- Murray, M. G. and Thompson, W. F. (1980) Rapid isolation of high molecular weight plant DNA. *Nucleic Acid Research* 8: 4321-4325.
- Muthusamy, S., Kanagarajan, S. and Ponnusamy, S. (2008) Efficiency of RAPD and ISSR markers system in accessing genetic variation of rice bean (*Vigna umbellata*) landraces. *Electronic Journal of Biotechnology* 11(3).
- Nkongolo, K. K., Michael, P. and Demers, T. (2005) Application of ISSR, RAPD, and cytological markers to the certification of *Picea mariana*, *P. glauca*, and *P. engelmannii* trees, and their putative hybrids. *Genome* 48: 302-311.
- Ozden-Tokatli, Y., Akdemir, H., Tilkat, E. and Onay, A. (2010) Current status and conservation of *Pistacia*

- germplasm. *Biotechnology Advances* 28: 130-141.
- Pirsevedi, S. M., Valizadehghan, S., Mardi, M., Ghaffari, M. R., Mahmoodi, P., Zahravi, M., Zeinalabedini, M. and Khayam Nekoui, S. M. (2010) Isolation and Characterization of Novel Microsatellite Markers in Pomegranate (*Punica granatum* L.). *International Journal of Molecular Science* 11: 2010-2016.
- Qian, W., Ge, S. and Hong, D. Y. (2001) Genetic variation within and among populations of a wild rice *oryza granulate* from China detected by RAPD and ISSR. *Theoretical and Applied Genetics* 102: 440-449.
- Rahimi, T., Sayed Tabatabaei, B. E., Sharifnabi, B. and Ghobadi, C. (2006) Genetic relationships between Iranian pomegranate (*Punica granatum* L.) cultivars, using Amplified Fragment Length Polymorphism (AFLP) marker. *Iranian Journal of Agricultural Science* 36: 1373-1379.
- Raina, S. N., Rani, V., Kojima, T., Ogihara, Y., Singh, K. P. and Devarumath, R. M. (2001) RAPD and ISSR fingerprints as useful genetic markers for analysis of genetic diversity, varietal identification, and phylogenetic relationships in peanut (*Arachis hypogaea*) cultivars and wild species. *Genome* 44: 763-772.
- Rohlf, F. J. (1998) NTSYS-PC Numerical Taxonomy and Multivariate Analysis System, Version 2.02. Exter Publications, Setauket, New York.
- Sarkhosh, A., Zamani, Z., Fatahi, R. and Ranjbar, H. (2009) Evaluation of genetic diversity among Iranian soft-seed pomegranate accessions by fruit characteristics and RAPD markers. *Scientia Horticulturae* 121: 313-319
- Smith, J. S. C., Chin, E. C. L., Shu, H., Smith, O.S., Wall, S. J., Senior, M. L., Mitchell, S. E., Kresovich, S. and Zeigle, J. (1997) An evaluation of the utility of SSR loci as molecular markers in maize (*Zea mays* L.): Comparison with data from RFLPs and pedigree. *Theoretical and Applied Genetics* 95: 163-173.
- Takeda, T., Shimada, T., Nomura, K., Ozaki, T., Haji, T., Yamaguchi, M. and Yoshida, M. (1998) Classification of apricot varieties by RAPD analysis. *Journal of Japanese Society of Horticultural Science* 67: 21-27.
- Talebi Bedaf, M., Ghobadi, C., Yamchi, A., Sayed Tabatabaei, B. E. and Bahar, M. (2005) Using of ISSR markers to investigate the genetic diversity in some Iranian pomegranate. *Proceedings of the 8th Congress of Agronomy and Plant breeding, Iran*.
- Wang, Z., Weber, J. L., Zhong, G. and Tanksley, S. D. (1994) Survey of plant short tandem DNA repeats. *Theoretical and Applied Genetics* 88: 1-6.
- Weiguo, Z., Zhihua, Z., Xuexia, M., Yong, Z., Sibao, W., Jianhua, H., Hui, X., Yile, P. and Yongping, H. (2007) A comparison of genetic variation among wild and cultivated *Morus* species (Moraceae: *Morus*) as revealed by ISSR and SSR markers. *Biodiversity and Conservation* 16: 275-290.
- Williams, J. G. K., Kubelik, A. E., Livak, K. J., Rafalski, J. A. and Tingey, S. C. (1990) DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research* 18: 6531-6535.
- Wunsch, A. and Hormaza, J. I. (2002) Cultivar identification and genetic fingerprinting of temperate fruit tree species using DNA markers. *Euphytica* 125: 59-67.
- Zamani, Z., Sarkhosh, A., Fatahi, R. and Ebadi, A. (2007) Genetic relationships among pomegranate genotypes studied by fruit characteristics and RAPD markers. *Journal of Horticultural Science and Biotechnology* 82: 11-18.
- Zietkiewicz, E., Rafalski, A. and Labuda, D. (1994) Genome fingerprinting by simple sequence repeat (SSR) anchored polymerase chain reaction amplification. *Genomics* 20: 176-183.

ارزیابی تنوع ژنتیکی ارقام انار ایران (*Punica granatum* L.) با استفاده از نشانگرهای RAPD و ISSR

مجید طالبی بدف^{۱*}، مسعود بهار^۱، بهرام شریف نبی^۱ و احد یامچی^۲

^۱ گروه بیوتکنولوژی، دانشکده کشاورزی، دانشگاه صنعتی اصفهان، اصفهان، ایران

^۲ دانشگاه علوم کشاورزی و منابع طبیعی گرگان

چکیده

با توجه به تنوع بالای مورفولوژیکی در ارقام انار ایران، تنوع ژنتیکی ۲۴ رقم انار با استفاده از نشانگرهای RAPD و ISSR مورد بررسی قرار گرفت. آغازگرهای RAPD در مجموع ۱۳۱ قطعه DNA تکثیر نمودند که ۲۹ قطعه آنها (۲۲/۱۴ درصد) چندشکل بودند. آغازگرهای ISSR نیز از مجموع ۱۷۳ قطعه تکثیر شده، ۲۹ (۳۷ درصد) چندشکلی حاصل نمودند. میانگین محتوای اطلاعات چندشکل (PIC) برای آغازگرهای RAPD و ISSR به ترتیب ۰/۱۲۸ و ۰/۱۶۳ به دست آمد. نتایج نشان داد که نشانگرهای ISSR در مقایسه با نشانگرهای RAPD الگوی نواری تکرارپذیری ایجاد می کنند و برای گروه بندی ارقام انار مؤثرترند. ضریب شباهت بین ارقام از ۰/۳۵۳ تا یک (RAPD) و ۰/۲۹۱ تا ۰/۹۳۰ (ISSR) متغیر بود و میانگین آن در نشانگرهای RAPD و ISSR به ترتیب ۰/۶۰۴ و ۰/۶۴۷ گزارش شد. تجزیه واریانس مولکولی (AMOVA) در داده های RAPD و ISSR اختلاف معنی داری بین نواحی مختلف جغرافیایی و طعم میوه ارقام مورد مطالعه نشان نداد ($P > 0.05$) که بیانگر عدم ارتباط تنوع جغرافیایی و تنوع ژنتیکی است.

واژه های کلیدی: *Punica granatum*، تنوع ژنتیکی، انار، RAPD، ISSR